



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

When $p=r/s$, $2m/(p^2-2)=2ms^2/(r^2-2s^2).....(3)$.

When $p=2(r+s)/(r+2s)$, $2m/(p^2-2)=m(r+2s)^2(r^2-2s^2).....(4)$.

Since (3) is integral, (4) is also.

Also solved by the *PROPOSER*.

GEOMETRY.

200. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

Find the locus of eight points of contact of the four common tangents of two concentric coaxial ellipses.

Solution by G. B. M. ZERR, A. M., Ph.D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let $x^2/a^2 + y^2/b^2 = 1$, and $x^2/c^2 + y^2/d^2 = 1$, be the equations to the ellipses, $c > a$, $b > d$.

Since $xx_1/a^2 + yy_1/b^2 = 1$, and $xx_2/c^2 + yy_2/d^2 = 1$, are two equations for the same line, we get $x_2 = c^2 x_1/a^2$, $y_2 = d^2 y_1/b^2$.

$\therefore x_1^2/a^2 + y_1^2/b^2 = 1$, and $c^2 x_1^2/a^4 + d^2 y_1^2/b^4 = 1$, give

$$x_1 = \frac{a^2 \sqrt{[b^2 - d^2]}}{\sqrt{[b^2 c^2 - a^2 d^2]}} = a^2 m \text{ (suppose)}, y_1 = \frac{b^2 \sqrt{[c^2 - a^2]}}{\sqrt{[b^2 c^2 - a^2 d^2]}} = b^2 n,$$

$$x_2 = c^2 m, y_2 = d^2 n.$$

Hence $mx \pm ny \pm 1 = 0$ represents the four common tangents. While $(c^2 m, d^2 n)$; $(a^2 m, b^2 n)$; $(-a^2 m, b^2 n)$; $(-c^2 m, d^2 n)$; $(-c^2 m, -d^2 n)$; $(a^2 m, -b^2 n)$; $(c^2 m, -d^2 n)$ are the eight points of contact. These eight points are situated on an ellipse, which is the locus required.

Let $x^2/A^2 + y^2/B^2 = 1$ be this ellipse. Then $c^2 m/A^2 + d^4 n^2/B^2 = 1$, also $a^4 m^2/A^2 + b^4 n^2/B^2 = 1$.

$$\therefore A^2 = \frac{(b^4 c^4 - a^4 d^4)m^2}{b^4 - d^4} = \frac{b^2 c^2 + a^2 d^2}{b^2 + d^2}, B^2 = \frac{(b^4 c^4 - a^4 d^4)n^2}{c^4 - a^4} = \frac{b^2 c^2 + a^2 d^2}{a^2 + c^2}.$$

$$\therefore (b^2 + d^2)x^2 + (a^2 + c^2)y^2 = b^2 c^2 + a^2 d^2 \text{ is the locus.}$$

201. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

Two plane sections of a right circular cone have their major axes AA' , aa' coplanar, and Aa on one generator equal to $A'a'$ on the other. The projections of the sections on any plane perpendicular to the axis are confocal.